

[ExPASy Home page](#)[Site Map](#)[Search ExPASy](#)[Contact us](#)[Swiss-Prot](#)

Hosted by NCSC US

Mirror sites:

[Canada](#)[China](#)[Korea](#)[Switzerland](#)[Taiwan](#)

Search

Swiss-Prot/TrEMBL



for

Go

Clear

# NiceProt View of Swiss-Prot: Q9BYT8

[Printer-friendly view](#)[Quick BlastP search](#)

[\[General\]](#) [\[Name and origin\]](#) [\[References\]](#) [\[Comments\]](#) [\[Cross-references\]](#) [\[Keywords\]](#) [\[Features\]](#)  
[\[Sequence\]](#) [\[Tools\]](#)

*Note: most headings are clickable, even if they don't appear as links. They link to the user manual or other documents.*

## General information about the entry

Entry name	NEUL_HUMAN
Primary accession number	Q9BYT8
Secondary accession number	Q9ULJ4
Entered in Swiss-Prot in	Release 41, February 2003
Sequence was last modified in	Release 41, February 2003
Annotations were last modified in	Release 42, September 2003

## Name and origin of the protein

Protein name	Neurolysin, mitochondrial [Precursor]
Synonyms	EC <a href="#">3.4.24.16</a> Neurotensin endopeptidase Mitochondrial oligopeptidase M Microsomal endopeptidase MEP
Gene name	NLN or KIAA1226
From	<a href="#">Homo sapiens (Human)</a> [TaxID: <a href="#">9606</a> ]
Taxonomy	<a href="#">Eukaryota</a> ; <a href="#">Metazoa</a> ; <a href="#">Chordata</a> ; <a href="#">Craniata</a> ; <a href="#">Vertebrata</a> ; <a href="#">Euteleostomi</a> ; <a href="#">Mammalia</a> ; <a href="#">Eutheria</a> ; <a href="#">Primates</a> ; <a href="#">Catarrhini</a> ; <a href="#">Hominidae</a> ; <a href="#">Homo</a> .

## References

[1]	SEQUENCE FROM NUCLEIC ACID. <a href="#">Chen J.M.</a> , <a href="#">Rawlings N.D.</a> , <a href="#">Barrett A.J.</a> ; "Cloning and sequencing of human neurolysin, an oligopeptidase of family M3."; Submitted (JAN-2001) to the EMBL/GenBank/DDBJ databases.
[2]	SEQUENCE FROM NUCLEIC ACID. <a href="#">TISSUE=Brain</a> ; MEDLINE=20039619; PubMed=10574462; [ <a href="#">NCBI</a> , <a href="#">ExPASy</a> , <a href="#">EBI</a> , <a href="#">Israel</a> , <a href="#">Japan</a> ] <a href="#">Nagase T.</a> , <a href="#">Ishikawa K.-I.</a> , <a href="#">Kikuno R.</a> , <a href="#">Hirosawa M.</a> , <a href="#">Nomura N.</a> , <a href="#">Ohara O.</a> ; "Prediction of the coding sequences of unidentified human genes. XV. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro."; DNA Res. 6:337-345(1999).

## Comments

- **FUNCTION:** Hydrolyzes oligopeptides such as neurotensin, bradykinin, dynorphin A, etc. (*By similarity*).
- **CATALYTIC ACTIVITY:** Preferential cleavage in neurotensin: 10-Pro<sup>-</sup>-Tyr-11.
- **COFACTOR:** BINDS 1 ZINC ION (*By similarity*).
- **SUBCELLULAR LOCATION:** MITOCHONDRIAL INTERMEMBRANE SPACE AND ALSO CYTOPLASMIC (*By similarity*).
- **SIMILARITY:** BELONGS TO PEPTIDASE FAMILY M3.

### Copyright

This SWISS-PROT entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to [license@isb-sib.ch](mailto:license@isb-sib.ch))

### Cross-references

EMBL	AJ300837; CAC27329.1; -. [ <a href="#">EMBL</a> / <a href="#">GenBank</a> / <a href="#">DDBJ</a> ] [ <a href="#">CoDingSequence</a> ] AB033052; BAA86540.2; -. [ <a href="#">EMBL</a> / <a href="#">GenBank</a> / <a href="#">DDBJ</a> ] [ <a href="#">CoDingSequence</a> ]
Genew	HGNC:16058; NLN.
CleanEx	HGNC:16058; NLN.
Ensembl	Q9BYT8; Homo sapiens. [ <a href="#">Entry</a> / <a href="#">Contig view</a> ]
HUGE	KIAA1226.
InterPro	IPR001567; Peptidase_M3. IPR006025; Zn_MTpeptdse. <a href="#">Graphical view of domain structure</a> .
Pfam	PF01432; Peptidase_M3; 1.
PROSITE	PS00142; ZINC_PROTEASE; 1.
ProDom	[ <a href="#">Domain structure</a> / <a href="#">List of seq. sharing at least 1 domain</a> ].
BLOCKS	Q9BYT8.
ProtoNet	Q9BYT8.
ProtoMap	Q9BYT8.
PRESAGE	Q9BYT8.
DIP	Q9BYT8.
ModBase	Q9BYT8.
SWISS-2DPAGE	<a href="#">Get region on 2D PAGE</a> .

### Keywords

**Metalloprotease; Hydrolase; Zinc; Mitochondrion; Transit peptide.**

### Features



[Feature table viewer](#)

Key	From	To	Length	Description
TRANSIT	1	37	37	MITOCHONDRION (BY SIMILARITY).
CHAIN	38	704	667	NEUROLYSIN.
METAL	497	497		ZINC (CATALYTIC) (BY SIMILARITY).
ACT_SITE	498	498		BY SIMILARITY.
METAL	501	501		ZINC (CATALYTIC) (BY SIMILARITY).
METAL	504	504		ZINC (CATALYTIC) (BY SIMILARITY).

**Sequence information**Length: **704 AA** [This is the length of the unprocessed precursor]Molecular weight: **80651 Da** [This is the MW of the unprocessed precursor]CRC64: **80136688D79BBEDF** [This is a checksum on the sequence]

10	20	30	40	50	60
MIARCLLAVR	SLRRVGSRI	LLRMTLGREV	MSPLQAMSSY	TVAGRNVLRW	DLSPEQIKTR
70	80	90	100	110	120
TEELIVQTKQ	VYDAVGMLGI	EEVTYENCLQ	ALADVEVKYI	VERTMLDFPQ	HVSSDKEVRA
130	140	150	160	170	180
ASTEADKRLS	RFDIEMSMRG	DIFERIVHLQ	ETCDLGKIKP	EARRYLEKSI	KMGKRNLHL
190	200	210	220	230	240
PEQVQNEIKS	MKKRMSELCI	DFNKNLNEDD	TFLVFSKAEI	GALPDDFIDS	LEKTDDDKYK
250	260	270	280	290	300
ITLKYPHYFP	VMKKCCIPET	RRRMEMAFNT	RCKEENTIIL	QQLPLRTRKV	AKLLGYSTHA
310	320	330	340	350	360
DFVLEMNTAK	STSRVTAFLD	DLSQKLKPLG	EAEREFILNL	KKKECKDRGF	EYDGKINAWD
370	380	390	400	410	420
LYYYMTQTEE	LKYSIDQEFL	KEYFPPIEVVT	EGLLNTYQEL	LGLSFEQMTD	AHVWNKSVTL
430	440	450	460	470	480
YTVKDKATGE	VLGQFYLDLY	PREGKYNHAA	CFGLQPGCLL	PDGSRMMAVA	ALVVNFSQPV
490	500	510	520	530	540
AGRPSLLRHD	EVRTYFHEFG	HVMHQICAQT	DFARFSGTNV	ETDFVEVPST	MLENWWVDVD
550	560	570	580	590	600
SLRRLSKHYK	DGSPHADDLL	EKLVASRLVN	TGLLTLRQIV	LSKVDQSLHT	NTSLDAASEY
610	620	630	640	650	660
AKYCSEILGV	AATPGTNMPA	TFGHLAGGYD	GQYYGYLWSE	VFSMDMFYSC	FKKEGIMNPE
670	680	690	700		
VGMKYRNLIL	KPGGSLDGMD	MLHNFLKREP	NQKAFILMSRG	LHAP	

Q9BYT8 in FASTA  
format[View entry in original Swiss-Prot format](#)[View entry in raw text format \(no links\)](#)[Report form for errors/updates in this Swiss-Prot entry](#)



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books	
Search	PubMed	▼	for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details			

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

☐ 1: Br J Pharmacol 1997 Jun;121(4):705-10

[Related Articles, Links](#)

Entrez  
PubMed

Full text article at  
[www.brjpharmacol.org](http://www.brjpharmacol.org)

### **Effect of a novel selective and potent phosphinic peptide inhibitor of endopeptidase 3.4.24.16 on neurotensin-induced analgesia and neuronal inactivation.**

PubMed  
Services

**Vincent B, Jiracek J, Noble F, Loog M, Roques B, Dive V, Vincent JP, Checler F.**

IPMC du CNRS, UPR411, Valbonne, France.

Related  
Resources

1. We have examined a series of novel phosphinic peptides as putative potent and selective inhibitors of endopeptidase 3.4.24.16. 2. The most selective inhibitor, Pro-Phe-psi(PO<sub>2</sub>CH<sub>2</sub>)-Leu-Pro-NH<sub>2</sub> displayed a K<sub>i</sub> value of 12 nM towards endopeptidase 3.4.24.16 and was 5540 fold less potent on its related peptidase endopeptidase 3.4.24.15. Furthermore, this inhibitor was 12.5 less potent on angiotensin-converting enzyme and was unable to block endopeptidase 3.4.24.11, aminopeptidases B and M, dipeptidylaminopeptidase IV and proline endopeptidase. 3. The effect of Pro-Phe-psi(PO<sub>2</sub>CH<sub>2</sub>)-Leu-Pro-NH<sub>2</sub>, in vitro and in vivo, on neurotensin metabolism in the central nervous system was examined. 4. Pro-Phe-psi(PO<sub>2</sub>CHH<sub>2</sub>)-Leu-Pro-NH<sub>2</sub> dose-dependently inhibited the formation of neurotensin 1-10 and concomittantly protected neurotensin from degradation by primary cultured neurones from mouse embryos. 5. Intracerebroventricular administration of Pro-Phe-psi(PO<sub>2</sub>CH<sub>2</sub>)-Leu-Pro-NH<sub>2</sub> significantly potentiated the neurotensin-induced antinociception of mice in the hot plate test. 6. Altogether, our study has established Pro-Phe-psi(PO<sub>2</sub>CH<sub>2</sub>)-Leu-Pro-NH<sub>2</sub> as a fully selective and highly potent inhibitor of endopeptidase 3.4.24.16 and demonstrates, for the first time, the contribution of this enzyme in the central metabolism of neurotensin.

PMID: 9208137 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	----------	---	------	---	---------	------	---

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)